


# How to use Auvi-Q<sup>®</sup> (epinephrine injection, USP)

## In case of an anaphylactic reaction:


Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.



**1** Pull off the RED safety guard.



**2** Place black end against outer thigh, then press firmly and hold for 5 seconds.




After using Auvi-Q, seek emergency medical attention immediately.

## Practicing with the Auvi-Q Trainer

Practice with the Auvi-Q Trainer before an allergic emergency happens to make sure you are able to safely use the active Auvi-Q in an emergency. After practicing, replace the RED safety guard and slide the trainer back into the gray outer case to reset.

**NOTE:** Every Auvi-Q prescription contains a trainer for practicing, along with 2 active devices.



Scan this QR code to watch a demo video.



## Indication

Auvi-Q<sup>®</sup> (epinephrine injection, USP) is used to treat life-threatening allergic reactions (anaphylaxis) in people who are at risk for or have a history of these reactions.

## Important Safety Information

Auvi-Q is for immediate self (or caregiver) administration and does not take the place of emergency medical care. Seek immediate medical treatment after use. Each Auvi-Q contains a single dose of epinephrine. **Auvi-Q should only be injected into your outer thigh. DO NOT INJECT INTO BUTTOCK OR INTRAVENOUSLY.** If you accidentally inject Auvi-Q into any other part of your body, seek immediate medical treatment. Epinephrine should be used with caution if you have heart disease or are taking certain medicines that can cause heart-related (cardiac) symptoms.

**If you take certain medicines, you may develop serious life-threatening side effects from epinephrine.** Be sure to tell your doctor all the medicines you take, especially medicines for asthma. Side effects may be increased in patients with certain medical conditions, or who take certain medicines. These include asthma, allergies, depression, thyroid disease, Parkinson's disease, diabetes, high blood pressure, and heart disease.

The most common side effects may include increase in heart rate, stronger or irregular heartbeat, sweating, nausea and vomiting, difficulty breathing, paleness, dizziness, weakness or shakiness, headache, apprehension, nervousness, or anxiety. These side effects go away quickly, especially if you rest.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please see accompanying full Prescribing Information or visit <http://products.sanofi.us/auvi-q/auvi-q.html>.